



Well Child/Immunization Schedule\*

Check-up Age	Immunizations
1 week old (2-4 days after hospital discharge)	Hep B (if not given in the hospital)
1 month	Hep B (if given at least 4 weeks apart)
2 months	DTaP, Hib, IPV, (Pentacel 3 in 1 combo) Prevnar 13, Rotateq (Oral) and Hep B (if not given at 1 month visit)
4 months	DTaP, Hib, IPV, (Pentacel 3 in 1 combo) Prevnar 13, Rotateq (Oral)
6 months	DTaP, Hib, IPV, (Pentacel 3 in 1 combo) Prevnar 13, Rotateq (Oral)
9 months	Hep B
12 months	MMR / VARIVAX, Hep A
15 months	DTaP, Hib, IPV, (Pentacel 3 in 1 combo) Prevnar 13, Rotateq (Oral)
18 months	DTaP, Hib, IPV, (Pentacel 3 in 1 combo) Prevnar 13, Rotateq (Oral)
Under 2 years	Hepatitis A ( <b>can now be given at 1 year of age</b> )
2 years	Prevnar 13 Booster / Hep A (2nd dose, if needed)
30 months	The AAP (American Academy of Pediatrics) recommends a 30 month check-up. Immunizations to be determined at time of visit.
3 years	Prevnar 13 Booster (if needed) / / Hep A (2nd dose, if needed)
4 or 5 years	Prevnar 13 Booster (if needed) / Hep A (if needed), DTaP, MMR, IPV and Varivax
6 years and older	Any required Booster vaccines
9 years and older	Gardasil®
11-12 years and older	Meningococcal , TDaP
11 years and every 5-10 years after	TDaP Booster
<b>When recommended by your provider the following vaccines are available certain times of the year</b>	
Spring and summer	Meningococcal and TDaP Booster, when applicable
Fall and winter	Influenza and Pneumovax

\* This is a typical immunization schedule. Due to many changes in immunizations, your child may not follow this exact timing for administration.